

# Evaluation of the Evidence for the Trauma and Fantasy Models of Dissociation

Constance J. Dalenberg  
California School of Professional Psychology at Alliant  
International University, San Diego

Bethany L. Brand  
Towson University

David H. Gleaves and Martin J. Dorahy  
University of Canterbury

Richard J. Loewenstein  
Sheppard Pratt Health System, Baltimore, Maryland, and  
University of Maryland School of Medicine, Baltimore

Etzel Cardëa  
Lund University

Paul A. Frewen  
University of Western Ontario

Eve B. Carlson  
National Center for Posttraumatic Stress Disorder, Menlo Park,  
and Veterans Administration Palo Alto Health Care System,  
Palo Alto, California

David Spiegel  
Stanford University School of Medicine

The relationship between a reported history of trauma and dissociative symptoms has been explained in 2 conflicting ways. Pathological dissociation has been conceptualized as a response to antecedent traumatic stress and/or severe psychological adversity. Others have proposed that dissociation makes individuals prone to fantasy, thereby engendering confabulated memories of trauma. We examine data related to a series of 8 contrasting predictions based on the trauma model and the fantasy model of dissociation. In keeping with the trauma model, the relationship between trauma and dissociation was consistent and moderate in strength, and remained significant when objective measures of trauma were used. Dissociation was temporally related to trauma and trauma treatment, and was predictive of trauma history when fantasy proneness was controlled. Dissociation was not reliably associated with suggestibility, nor was there evidence for the fantasy model prediction of greater inaccuracy of recovered memory. Instead, dissociation was positively related to a history of trauma memory recovery and negatively related to the more general measures of narrative cohesion. Research also supports the trauma theory of dissociation as a regulatory response to fear or other extreme emotion with measurable biological correlates. We conclude, on the basis of evidence related to these 8 predictions, that there is strong empirical support for the hypothesis that trauma causes dissociation, and that dissociation remains related to trauma history when fantasy proneness is controlled. We find little support for the hypothesis that the dissociation–trauma relationship is due to fantasy proneness or confabulated memories of trauma.

Keywords: trauma, dissociative disorder, dissociation, suggestibility, fantasy

Scientific interest in the concept of dissociation and the etiology researchers have empirically identified and investigated various of the dissociative disorders has increased markedly in recent types and categories of dissociation: the identity alterations and decades. Building on the foundational work of Janet (1889, 1919), amnesias prominent in the dissociative disorders (Putnam, 1991),

---

This article was published Online First March 12, 2012. Center for Posttraumatic Stress Disorder, Menlo Park, and Veterans Administration Palo Alto Health Care System, Palo Alto, California; David H. Gleaves is now at School of Psychology, Social Work and Social Policy, University of South Australia, Adelaide, Australia. We thank Franziska Unholzer for assistance with constructing Table 5. Correspondence concerning this article should be addressed to David Spiegel, Department of Psychiatry and Behavioral Sciences, Stanford University, 401 Quarry Road, Office 2325, Stanford, CA 94305-5718. E-mail: dspiegel@stanford.edu



ported trauma and dissociation, but ascribe different reasons for the relationship. It is therefore important to clarify the true points of distinction in the two models. These appear to fall into eight

and suggestibility, and therefore predicts little to no relationship between dissociation and trauma if fantasy proneness and suggestibility are controlled. Alternatively, the TM clearly predicts an increment for trauma over fantasy proneness or suggestibility. In contrast, the causal path for the FM does not posit a role (or at least a significant role) for trauma in the neuropsychological or cognitive deficits seen in dissociative individuals. In Merckelbach et al.'s (2002) model, for instance, the relationship between dissociation and trauma self-report was hypothesized to be fully mediated by absent-mindedness and fantasy proneness, with no role for actual trauma. Cognitive deficiencies inherent to dissociation were thought to be a primary source of the trauma report.

### **Predictions Regarding Omission and Fragmentation of Memory**

The TM posits that the dissociative individual is largely attempting to avoid recall of trauma by conscious and unconscious disavowal of the importance, implications, and/or accuracy or reality of the memory. According to the TM, the dissociative individual attempts to avoid thinking about the memory, disconnects from the emotional content of the memory, and ultimately may fail to recall some or all of the memory (e.g., DePrince & Freyd, 2004; Dorahy, 2006). The avoidance associated with dissociation may be both conscious and unconscious, or may be an initially conscious process that becomes unconscious over time (see Erdelyi, 1990). Automatic withdrawal of attention upon exposure to trauma or reminders of trauma, potentially resulting from dissociative episodes during encoding, may inhibit associative processing (Lyttle, Dorahy, Hanna, & Huntjens, 2010), and may result in a lack of the rich associative network typical of important emotional memories (cf. Spiegel & Cardena, 1991; Stern, 1997). The result is a set of nonintegrated and fragmented memories (data driven/perceptual rather than autobiographical/conceptual; Brewin, Dalgleish, & Joseph, 1996; Ehlers & Clark, 2000; Holmes, Brewin, & Hennessey, 2004). This type of processing might account for omissions and poor agreement in detail across narrative recountings. Over the course of time, fragmented memories lacking associative networks may be more easily forgotten. This reasoning supports TM hypotheses regarding relationships between dissociation and fragmentation of memory and between dissociation and lost or recovered memory.

FM theorists make no claim for the relationship of fragmentation and dissociation. Omission, however, is thought by FM theorists to be negatively related to dissociation (cf. Giesbrecht et al., 2008). The FM argument here is that any elevation in trauma report by dissociative individuals is due to exaggeration and fantasy. Therefore, omission of data and loss of detail in severe trauma is less likely for dissociative individuals than is addition of detail and enhancement of the trauma description.

### **Predictions Regarding the Biology and Neurobiology of Trauma**

Both the TM and the FM are consistent with a biological or sociobiological foundation for dissociation. The TM predicts that the experience of trauma and high levels of stress are related to cognitive deficits (Vasterling et al., 2002). The effects will appear in individuals with clinical dissociative disorders, as well as in traumatized nondissociative individuals, and will include the errors of omission, commission, and narrative fragmentation mentioned earlier (Harvey & Bryant, 1999; Kleim, Wallott, & Ehlers, 2008). Further, TM theorists expect differences between dissociative and nondissociative individuals in neurobiological studies, such as in

sion and increased fragmentation. The FM, which presents dissociation as related to exaggeration and false generation of trauma (Dell, 2006; see Carlson, 2008, for a review of the predicts no relationship or a negative relationship between dissociation and fragmentation or omission).

7. The TM predicts that, over time, dissociative individuals will be more likely to "forget" or have difficulty accessing important facets of the memory. The FM states that those who claim recovery of a memory are unlikely to be recalling an actual trauma.

8. Both models predict some relationship between dissociation and neuropsychological measures such as working memory (similar to those seen in work with PTSD; Vasterling & Brewin, 2005). The TM holds that the biology of dissociation will ultimately fit with a theory of a brain-based regulatory response to fear or other extreme emotion (Lanius et al., 2010). Thus, the psychophysiology of the dissociative individual should be differentiable from the nondissociative individual in fear-relevant situations. The FM makes no prediction in this area.

### Measurement of Dissociation and Fantasy Proneness

Prior to the analysis of the evidence for the TM and FM of dissociation, attention should be given to the measurement of this construct. The DES (Bernstein & Putnam, 1986) has been used in over 2,000 studies of dissociation to date, as both the focus for reviews of positive findings and the central instrument cited by critics of dissociation and its measurement. The DES also has an adolescent variant (the Adolescent Dissociative Experiences Scale [ADES]; Armstrong, Putnam, Carlson, Libero, & Smith, 1997) and a checklist form for use by parents or other adults assessing young children (Child Dissociative Checklist [CDC]; Putnam, Helmers, & Trickett, 1993).

In addition to the DES and its variants, a number of alternative instruments have appeared, such as the Questionnaire of Experiences of Dissociation (Riley, 1988) and the Dissociation Questionnaire (Vanderlinden, Van Dyck, Vandereycken, Vertommen, & Verkes, 1993), but these alternatives have not received substantial research attention. Briere's (2002) Multiscale Dissociation Inventory (MDI) is a promising new addition to the library of dissociation measures, particularly given the availability of clinical norms, but again little is yet available to establish the ability of the measure to tap important dissociation-related phenomena.

Wright and Loftus (1999) have developed a creative alternative to the DES. Using the same items as the DES, Wright and Loftus's DES-C asks participants not to rate their dissociative symptoms, but instead to rate whether they are dissociating less or more than others whom they know. The contention that this capacity is within the skill set of the dissociative patient (or even the normal control) has yet to be demonstrated. Further, the DES-C correlates only .25 with the DES (Wright & Loftus, 1999), clearly raising questions about the similarity of the two measures. We could find no published evidence showing that the DES-C is in fact a measure of dissociation. In the review below, research focuses on the original DES and its child and adolescent variants.

In addition, several diagnostic inventories and interviews have been developed for the diagnosis of clinical dissociative disorders. They are not discussed in detail here. However, they include two diagnostic interviews, the Structured Clinical Interview for DSM-IV–TR Dissociative Disorders (SCID-D; Steinberg, 1994) and the

### The Dissociative Experiences Scale

The DES is a 28-item self-report measure. In the original Bernstein and Putnam (1986) measure, the frequency of each item was rated along an 11-point visual analog scale. In a revision by E. B. Carlson and Putnam (1993), the scale was changed to a Likert model with choices ranging from 0% (never) to 100% (always) at 10 percentage point increments. A sample item is "Some people have the experience of finding themselves in a place and having no idea how they got there" (Item 3). The DES has also been shown to measure both a taxon, often described as "pathological" dissociation, typically measured by the eight-item dissociative taxon, or DES-T (Waller, Putnam, & Carlson, 1996), and a continuum, measured by the total scale or by the "nonpathological" absorption subscale (Waller et al., 1996). The DES-T consists of lower base rate items targeting measurement of depersonalization and derealization, identity fragmentation, and amnesia. The absorption subscale is a subset of higher base rate DES items assessing normal experiences of deep focal attention as well as lapses in attention.

Critics of the current measurement of dissociation and, in particular, of the DES tend to focus on three issues: the inclusion of absorption in the domain of dissociation, the reliability and meaning of the taxon, and the more general issue of giving a unitary label (dissociation) to a wide range of topics, often symbolized by the argument of whether the DES is unifactorial or multifactorial (Bernstein, Ellason, Ross, & Vanderlinden, 2001; Giesbrecht et al., 2008; Watson, 2003). The argument against the inclusion of absorption in the measurement of dissociation can be made in two ways: (a) that high absorption is not a symptom of dissociative disorders, because it is more common in the general population than DES taxon items, and (b) that absorption is normal and nonpathological at all levels. The first assumption is not supported by the empirical evidence. For example, approximately 75% of patients with diagnosed dissociative disorders in Leavitt's (2001) sample had high scores on absorption scales. Dalenberg and Paulson (2009), using a version of the DES corrected for skewness, found that over 95% of taxon-positive individuals were above the cutoff for high absorption. Further, the correlation between the taxon and absorption factors is very high ( $r = .80$ ; Levin & Spei, 2004;  $r = .36-.72$  in six psychiatric groups in Leavitt, 1999). These findings call into question the contention that "cleaner" measures of dissociation should exclude absorption. Instead, the data support the inclusion of items that measure capacities that may be facilitators, precursors, or lower level symptoms of dissociation.

With reference to the second assumption, high absorption has been shown repeatedly to be a marker for severe psychopathology.

Indeed, Allen, Coyne, and Console (1997) reported surprise that the nonpathological absorption facets of dissociation were more related to psychosis than were the taxonic items. Absorption correlated more highly with severe psychopathology on the Minnesota Multiphasic Personality Inventory and the Millon Multiaxial Inventory than did the amnesia and depersonalization factors (Allen et al., 2002).

TM theorists and FM theorists both share the concern that the DES-T yields unacceptably high false-positive rates if used as a sole diagnostic instrument (cf. Carlen 2008), and that it has modest reliability in nonclinical samples when dissociative disorder should be rare or nonexistent (.62 over 2 months; Watson,



Table 1  
Relationship of Trauma and Dissociation

Study	Participants	Trauma type	Trauma measure	Dissociation measure	r
Akyüz et al., 2005	251 adult women See above See above	Nonclinical samples PA SA TOT	CANQ	DES (Turkey)	.06
			CANQ	DES	.18
			CANQ	DES	.22
			UCLA-PTSD Index	DES	.34
Chu & DePrince, 2006 Collin-Vézina & Hebert, 2005	67 adult mothers 67 children evaluated for abuse and matched controls	SA	Hospital evaluation and interview	CDC	.38***
			Guardian report on UCLA-PTSD Index Guardian report on UCLA-PTSD Index	ADES CDC	.21 .25
Dorahy et al., 2007 Dutra et al., 2009	97 children 66 adults 56 young adults 56 young adults	TOT DV DMC TOT	TEC	DES	.40
			CTES-R	DES	.38*
			Behavioral codes on AMBIAC	DES	.16
			Self-report	DES (Netherlands)	.31
Geraerts et al., 2005 Kisiel & Lyons, 2001	98 adults 114 wards of DCFS 114 wards of DCFS 114 wards of DCFS	SA PA PA SA	Reported by DCFS caretaker	ADES	.20*
			Reported by DCFS caretaker	ADES	.24*
			Reported by DCFS caretaker	CDC	.32***
			Reported by DCFS caretaker	CDC	.30*
Macfie et al., 2001a Macfie et al., 2001b McNally et al., 2000 McNally et al., 2006	198 children 88 children 68 adults 166 adults	SA TR SA SA	CPS records	CDC	.11
			CPS records	CDC	.39***
			Coded from CPS records	CDC	.42***
			Self-report	DES	.32
Narang & Contreras, 2005 Nåring & Nijenhuis, 2005 Nilsson & Svedin, 2006 Noll et al., 2003 Ogawa et al., 1997	76 mothers 147 adults 391 adolescents 166 children 168 young adults	PA TOT TOT SA TR	CHQ	DES	.42*
			TEC	DES (Netherlands)	.37
			DIS-Q	DES (Sweden)	.27
			Substantiated by CPS	ADES	.28
Sayar et al., 2005 Smith et al., 2010 Somer, 2002 Trickett et al., 2001	173 adolescents 50 adults 90 adults 166 children	PA TOT TOT SA	Home observation, CPS records, parent interview	DES	.36***
			Self-report	ADES (Turkey)	.31
			THS	DES	.44
			TEQ	DES (Israel)	.39
Twaite & Rodriguez-Srednicki, 2004 Zoroglu et al., 2003	158 young adults 284 adults 839 adolescents	SA PA SA TR	Verified through DCFS	CDC at 6 months (Time 1)	.34***
			Verified through DCFS	ADES at 7 years after Time 1	.16*
			Self-report	DES	.24
			Self-report	DES	.18
Brunner et al., 2000 E. B. Carlson et al., 2001 Dell, 2006	198 adolescent inpatients See above 178 adult inpatients See above See above See above 204 clinical and nonclinical See above See above	Clinical samples SA PA Violent SA Other SA Violent PA Other PA PA SA TOT	Therapist reports based on guardian reports, and self-report	ADES (Germany)	.36***
			DCFS records, and self-report	ADES	.22***
			Structured interview	DES	.52
			See above	DES	.49
Dell, 2006	See above See above See above See above	Violent PA Other PA Other PA PA SA TOT	See above	DES	.35
			See above	DES	.28
			TEQ	DES	.34
			TEQ	DES	.44
			TEQ	DES	.47

(table continues)



Table 1 (continued)

Study	Participants	Trauma type	Trauma measure	Dissociation measure	r
El-Hage et al., 2002	140 adult outpatients	TR	CAPS	DES (France)	***.49
Francia-Martinez et al., 2003	100 adult inpatients	SA	BSAE	DES (Puerto Rico)	.14
Freyd et al., 2005	99 adults with chronic illness or pain	BT	BBTS	DES	.43**
Gast et al., 2001	102 adult inpatients	PA	CTQ	DES (Germany)	***.47

with no trauma of the type studied were included; (c) sample size was 50 or greater; and (d) the study used a community sample or a clinical sample including a range of psychiatric diagnoses. Thus, samples consisting entirely of dissociated patients or those with PTSD, which may have restricted values on trauma likelihood or dissociation, were not included, but consecutive psychiatric admissions samples of children in therapy are represented. College samples which are likely to be biased in favor of low impairment, were experienced (i.e., one trauma vs. no trauma  $Hedges'g = 0.56$ , not included. Lev-Wiesel, Daphna-Tekoah, and Hallak's (2009) two traumas vs. no trauma  $Hedges'g = 0.84$ , and three traumas large sample of pregnant women was not included given the complex literature on the relationship of pain, stress, and dissociation (cf. Ludacher et al., 2007). Studies using only certain physical abuse subscales of the DES also were not included. Studies that appeared to test the same sample in different publications and studies that limited trauma effects to emotional abuse were excluded. The effect size was chosen, since the majority of studies reported this figure.

Table 1 presents the results of 38 studies that met our criteria. The average weighted effect size was .31 for the 19 sexual abuse samples, .27 for the 12 physical abuse studies, and .34 for the 16 total trauma score studies (for the E. B. Carlson et al., 2001, study, the two relevant values were averaged). The overall weighted  $r$  estimate was .32. Fixed-point estimates were made via Comprehensive Meta-Analysis software. All values are moderate effect size  $Q$  values were between 24.59 (for sexual abuse) and 63.71 (for all studies), indicating heterogeneity of effect sizes.

The table also illustrates that large population studies and well-controlled comparison studies do exist that test the relationship of

that would allow an effect size computation, presenting instead compared with the average weighted for the remaining physical abuse large samples of dissociative disorder or dissociative identity disorder studies of .26. The objective physical abuse analysis yielded an order (DID) patients. Trauma history was found in 50%–100% of nonsignificant  $Q$  value of 3.67 ( $p > .05$ ), with the remaining such individuals in all studies (with the exception of the Turkish analyses showing  $Q$  values at or greater than 23.33 ( $p < .01$ )). study by Sar, Akyüz, & Dogan, 2007). These results also support These results contradict the FM prediction, and go to the heart of the TM, but differ from the data in Table 1. For Table 1, all clinical the FM argument. If the trauma–dissociation relationship were samples showed a general relationship between trauma and level largely due to fantasy proneness and subsequent exaggeration of of dissociativity on the DES. However, the base rate of DID in trauma, clearly the relationship should be weaker when trauma is most clinical samples is low (1.3% in Ross's, 1991, nonclinical measured with greater objectivity. This argument has been made sample); thus, the correlation coefficient can be misleading. For explicitly in Giesbrecht et al.'s (2008) recent review. They were instance, in a large sample ( $n = 618$ ), Briere's (2006) correlation able to locate two studies with objective criteria, both with small of .11 between trauma and clinical elevation on the MDI accounts and nongeneralizable samples, noting that neither reached statis- for less than 2% of the variance. If the same results are translated into the language of binomial probability to make base rates more reasonable samples, all of which did support the TM hypothesis, were not visible, as Briere made possible through cross-tabulation charts, discussed by those authors. In this full review comparing studies the probability of clinical elevation in the MDI is 4 times greater with self-report to those using objective measures, studies with in the trauma-exposed compared with a nonexposed sample (8% self-report measures of trauma did not show a greater relationship vs. 2%). Further, the probability of a trauma history given into dissociation than those with objective measures. Again, these elevated MDI in this sample was 90%. Similarly, in the study with findings support the TM position, not the FM view. the weakest effect size in Table 2 (Set et al., 2007), the probability There have been no large-scale studies of the objective evidence of abuse within the dissociative disorder samples were still 2–4 for trauma reported by dissociative disordered patients including times higher than the rates within psychiatric controls. The heterogeneity of effect sizes are reflected in the very high values of the base rate of dissociative disorders. However, the smaller studies that have followed up on the evidence for child trauma history in DID patients have confirmed the existence of such trauma. sexual abuse. The mean-weighted was .54 for the five sexual abuse samples and .52 for the five physical abuse samples. Again, Coons (1994) found documented corroboration (e.g., CPS and with diagnosis rather than dissociation as a continuum, the hypothesis of the consistent relationship between trauma and dissociation is supported. Coons and Milstein (1986) found documentation through medical records or family testimony in 17 of 20 adult DID patients (see also Lewis, Yeager, Swica, Pincus, & Lewis, 1997). Further, Hornstein and Putnam (1992) described two samples of children and adolescents with dissociative disorders totaling 74 participants, all of whom had reported histories of a wide variety of types

## Evidence for Prediction 2: Does the Trauma–Dissociation Relationship Disappear in Studies With “Objective” Measures of Trauma?

In Table 1, 10 studies included external criteria for determination of maltreatment status, including physical abuse, sexual abuse, witnessing of maltreatment, parental death, and/or neglect. Social service investigation the hypotheses of this review and blind to the results of each substantiated 95% of these histories. study—made this judgment with 100% agreement. The “objective” data included confirmation by therapists (with access to guardians and Child Protective Services [CPS] reports), protective agency report determined by researchers, or, in the case of Dutton Bureau, Holmes, Lyubchik, and Lyons-Ruth (2009), observer behavior codes of mothers' treatment of their infants. In Dutra et al. (1997), disrupted maternal communication included ratings of sexual & Putnam, 2003; Trickett, Noll, Reiffman, & Putnam, 2001), or withdrawal, and fearful-disoriented behavior on the part of the mother. For example, in an ongoing, case-controlled longitudinal study of mother in the Ainsworth Strange Situation task. Nine of these 10 girls with a substantiated history of child sexual abuse (CSA; studies tested the correlation between dissociation and sexual abuse, whereas three also tested the correlation between dissociation and physical abuse. The FM prediction that objectively determined trauma would show lower correlations with dissociation than self-reported trauma thus could be tested by comparing the effect size of the objective studies with the studies using a standardized self-report measure or a single-item self-label of sexual abuse. Using a weighted mean effect size, the objective studies on sexual abuse had a weighted average of .30, whereas the self-report, standardized measure, or structured interview studies had a weighted average effect size of .32. The three objective measure studies on physical abuse had a weighted average of .30, com-

related with self-reported dissociation at age 19. Within the disorganized group, higher dissociation scores were found for the group that had experienced documented traumas in childhood and adolescence. Ogawa et al. also reported a statistically significant correlation between trauma (with both objective and parent self-report documentation) and dissociation at Time 1 (infancy), Time 4 (age 16–17), and Time 5 (age 19) with a sample of 168. In Diseth's (2006) smaller study of children who had experienced repeated and painful medical procedures ( $n = 42$ ), an objective trauma, dissociation in both adolescence and young adulthood correlated with number of hospitalizations ( $r = .59$  with the ADES and  $r = .79$  with the DES, 10 years later).

Many prospective studies follow at-risk samples in order to have realistic probability of finding traumatized individuals with varying symptom levels. Barring the random (and unethical) assignment of individuals to traumatizing conditions, the optimal deter-

Over longer time spans, the TM prediction would be that traumatized individuals would be temporarily elevated in dissociative content of the CEQ and the DES. They suggested:

symptoms as a group, and that these symptoms would diminish for most individuals over time as the trauma becomes more integrated into cognitive systems and trauma-related emotions (e.g., fear and anxiety) dissipate. In studies in which participants were followed after trauma—as in Cardarand Spiegel (1993); E. B. Carlson et al. (2011); Feeny, Zoellner, Fitzgibbons, and Foa (2000); and Feeny, Zoellner, and Foa (2000)—large and statistically significant drops in dissociative symptom severity occur over time with- out intervention in most individuals. Two to 10 days after trauma

Two CEQ items (i.e., “I often confuse fantasies with real memories” and “I sometimes feel that I have an out of body experience”) clearly overlap with some DES items (e.g., “not sure whether one has done something or only thought about it” and “feeling as though one’s body is not one’s own,” respectively). (p. 989)

Further, it is consistent with prior theory and research on fantasy proneness scales that trauma is one cause, although not the sole cause, of fantasy proneness. In early articles on the CEQ, Merckelbach et al. (2001) conceded there are different paths to fantasy proneness, including coping with childhood adversity: “Other fantasy-prones,” they wrote, “reported a heightened frequency of aversive childhood events. In these cases, a profound fantasy life may have become a means to cope with or escape from negative experiences” (p. 988). Rhue and Lynn (1987, p. 121), for instance, noted that fantasy-prone participants reported “greater frequency and severity of physical punishment, greater use of fantasy to block the pain of punishment, more thoughts of revenge toward the person who punished them, greater loneliness, and a preference for punishing their own children less severely” than those lower in fantasy proneness. Lynn and Rhue (1986) and S. C. Wilson and Barber (1983) also reported that fantasizers acknowledged more severe and more frequent childhood punishment. In keeping with the TM hypothesis of use of fantasy as escape, fantasy proneness is related to the five scales of the Childhood Trauma Questionnaire (Pekala et al., 1999–2000).

In summary, the increase in state dissociation after exposure to high stressors or traumatic events and trauma reminders is consistent with TM Prediction 3. Similarly, findings support the TM prediction of the short-term decrease in dissociation (relative to comparison groups) with trauma-relevant psychological or pharmacological treatment and the long-term decrease in dissociation over time. If dissociation were a stable outgrowth of fantasy proneness and mild neurocognitive disturbance (cf. Giesbrecht et al., 2008), such patterns would be much harder to explain. These findings clearly support TM Prediction 3, that dissociation is temporally related to trauma and trauma treatment.

#### Evidence for Prediction 4: Does Dissociation Show an Increment Over Fantasy Proneness in the Prediction of Trauma?

Both the TM and the FM predict a relationship between the measures typically used for the dissociation and fantasy proneness concepts, because both types of scales were developed from a theoretical base that included an etiological role for psychological absorption and trauma. Fantasy proneness is acknowledged to be the “close cousin” of absorption by Geraerts, Merckelbach, Jelicic, Smeets, and Van Heerden (2006, p. 1143). The authors of both studies that track these relationships over time.

Although the relationship of fantasy proneness and dissociation is not incompatible with either model, the FM does make a prediction of the relative relationship of these variables to trauma replicated in Figure 1, a statistical prediction can be made that fantasy proneness will produce an increment over dissociation in self-report. In the FM given by Merckelbach et al. (2002), and Bernstein & Putnam, 1986). It is easy for theoreticians from all perspectives to lose track of this history, reifying the scale totals and reporting as an independent and surprising finding that absorption correlates strongly with each measure.

Merckelbach et al.’s (2001) CEQ and S. C. Wilson and Barber’s (1983) ICMI do correlate with dissociation (Merckelbach et al., 2002; Pekala et al., 1999–2000; Rauschenberger & Lynn, 1995; Waldo & Merritt, 2000), but the reason for the correlation is unclear. Highly fantasy-prone individuals have been reported to be diagnosed with dissociative disorders more often than low- or medium-level fantasy-prone individuals (Rauschenberger & Lynn, 2002–2003). The inclusion of absorption within each scale type is the most obvious explanation. Merckelbach et al., the developer and Thomson, Keehn, and Gumpel (2009). Specifically, in each

case, fantasy proneness did relate to trauma history and dissociation, but trauma history did have an increment over fantasy proneness in the equation predicting the DES. Dissociation does relate to report of trauma history controlling for fantasy proneness.

Furthermore, the few studies on fantasy proneness in dissociative disordered samples do not indicate the strong elevations in fantasy proneness that would be expected if their trauma histories were entirely fantasized. Huntjens et al. (2006) found that DID patients scored higher on fantasy proneness than controls and nonclinical DID simulators. However, the DID mean score on CEQ (9.92) was very similar to means of male and female college students reported in Merckelbach et al.'s (2001) psychometric article on the CEQ ( $M = 9.2, SD = 4.4$ , and  $M = 8.7, SD = 4.0$ , respectively). Using the ICMI, Levin, Sirof, Simeon, and Guralnick (2004) also found elevated levels of fantasy proneness in patients with depersonalization disorder (DPD) compared with nonsymptomatic controls. However, as Levin et al. wrote, the scores for the DPD group were well below typically used thresholds for high fantasy proneness. The DPD mean was  $SD = 7.3$ ), which falls at the low end of the range for medium fantasy proneness on this instrument (14–36 in Levin et al., 2004). These findings in general support the TM prediction (Prediction 4) regarding the independent contribution of dissociation over fantasy proneness in the prediction of trauma history.

### Evidence for Prediction 5: Are Dissociative Research Participants at High Risk for Suggestibility and False Memory?

Research on suggestibility is also central to the FM contentions about the dissociation–trauma connection. The controversial contentions of the FM are not only that the dissociation and trauma report connection is mediated by fantasy proneness, which appears unfounded as discussed earlier, but also that dissociation produces enhanced probability of confabulation of trauma memory itself. Research testing general memory skills of dissociative individuals, as Giesbrecht et al. (2008) repeatedly cited their concern that dissociative individuals will overreport trauma on standardized questionnaires unless provided with a context that “discourages reporting of traumatic experiences” (p. 622). It seems ill-advised and potentially harmful to discourage patients from reporting trauma exposure due to fears of high rates of false report without strong support for this hypothesis.

**Suggestibility paradigms.** In the standard FM argument of dissociation as a risk factor for suggestibility, many nonequivalent forms of suggestibility are mentioned and tested (Giesbrecht et al., 2008; Merckelbach & Muris, 2001). To extend the range of studies reported, all research with samples greater than 25 are presented in Table 4. The best known are clustered under event suggestibility studies, and represent forms of suggestion that include acceptance of the false suggestion that one has seen or experienced an event. In the nonautobiographical studies of this type, participants are typically shown slides or read paragraphs, and pressed at a later point to agree to a false statement about a slide seen or fact heard. The Gudjonsson (1997) suggestibility paradigm is a standardized form of this type of suggestibility. In this paradigm, participants are read paragraphs and then (through social pressure or misleading questions) pushed toward acceptance of false statements about the information heard. An overall suggestibility score, a yield score (degree of acquiescence to leading questions), and a shift

score (the number of times the individual changed an answer in response to interpersonal pressure) are then calculated. The methodology in autobiographical event suggestibility studies is more varied. In studies typically referred to as “false memory” studies (e.g., Hyman & Billings, 1998), participants are told that a knowledgeable person (typically the individual’s mother) recalls an event in the person’s life. The dependent variable is the degree to which the research participant appears to accept the truth and this false memory. In misinformation studies, the dependent variable is the same, but the procedures typically involve less powerful suggestion (misleading questions, varying in terms of source, number, and strength). In source monitoring or source confusion studies, the task of the participant is typically to discriminate between competing sources (picture seen, a paragraph read, or a new story heard). Alternatively, in the Deese–Roediger–McDermott (DRM) paradigm (Deese, 1959; Roediger & McDermott, 1995), the participants read a series of words that relate to an overarching nonpresented word (e.g., read the words *sap, doze, and dream*—all words related to the concept “sleep”). The dependent variable is whether the individual recalls or falsely assents to seeing the nonpresented concept word.

Finally, in the imagination inflation studies, participants imagine a series of incidents and are asked about their feeling of remembering the event, as opposed to merely knowing or believing that the event might have happened. The events are typically plausible or known events from childhood.

The degree to which each of these paradigms is linked to a general “suggestibility” trait is unknown, but sets of studies are reviewed in turn as examples of suggestibility as defined within the FM. Historically, false memory has been fairly loosely defined in such paradigms (cf. DePrince, Allard, Oh, & Freyd, 2004). Research testing general memory skills of dissociative individuals, or errors on event memory tasks in the absence of suggestion, are not considered as examples of suggestibility paradigms.

**Nonautobiographical event suggestibility.** Table 4 contains data from eight studies with 10 samples investigating suggestibility for nonautobiographical events, all using the Gudjonsson methodology, and the examination of suggestibility relationship with dissociative experiences. The clinical samples—a small group of anxious patients reported by Wolfradt and Meyer (1998) and the larger mixed sample by Little (1996)—and the only abuse sample (Schultz, Passmore, & Yoder, 2003) produced nonsignificant results. The weighted estimate for the correlation between dissociation and suggestibility in this category is .12. Further, the pattern of correlations on the Gudjonsson subscales varied across the few studies reporting statistically significant results. Wolfradt and Meyer in their nonclinical sample found DES correlations with both Shift and Yield scales; Merckelbach, Muris, Rassin, and Horselenberg (2000) reported DES correlations with the Shift (but not Yield) score; and Merckelbach, Muris, Wessel, and Van Koppen (1998) found correlations with the Yield (but not Shift) score. Horselenberg et al. (2000) came to the conclusion that “the relationship between dissociative tendencies and memory distortions is not as impressive as some authors have suggested” (p. 136), noting that the few previous studies that had found positive associations had methodological limitations. Gudjonsson (2003) himself specifically noted with surprise the lack of consistent relationship

Table 4  
Relationship of Dissociation and Suggestibility

Study	Sample	Dissociation measure	Suggestibility task	r
Nonautobiographical event suggestibility				
Haraldsson, 2003	30 children, half with reincarnation beliefs	CDC	E; GSS	.11
Hekkanen & McEvoy, 2002	111 UG	DES	E; acceptance of false suggestions regarding slides	.00
Horselenberg et al., 2004	38 UG	DES	E; GSS	.08
Merckelbach, Murrin, Rassin, & Horselenberg, 2000	56 female UG	DES	S; GSS	** .37
Merckelbach et al., 1998	40 UG in Study 1	DES	S; GSS	* .32
Schultz et al., 2003	51 CM and 31 RM sexual abuse survivors	DES	E; GSS	-.06 and .06
Torrens, 2005	146 UG	DES	E; GSS	.06
Wolfraedt & Meyer, 1998	45 controls and 37 anxious patients	DES	E; GSS	* <del>.06</del> -.04
Autobiographical events				
Drivdahl & Zaragoza, 2001	149 UG	DES	E; errors after suggestion regarding staged event	.07
Eisen & Carlson, 1998	130 UG	DES	E; agreement with misleading information after staged event	** -.21
Eisen, Morgan, & Mickes, 2002	111 UG	DES	E; agreement with misleading information after staged event	.13
Eisen, Qin, et al., 2002	49 children	CDC	E; agreement with misleading information after abuse assessment	.01
Horselenberg et al., 2004	38 UG	DES	E; false recognition of foils regarding autobiographical events	.19





but also may occur under event suggestibility paradigms that convince the participant of the truth of a nonremembered event and encourage general attempts to remember (Hyman & Billings, 1998; Ost et al., 1997).

A challenge for suggestibility theorists is the differentiation of acquiescence and false memory. If the task is to remember an

(2007) found spontaneously recovered memories to be similarly likely to have corroboration (37%) when compared with continuous memories (corroborated in 45%). However, memories recovered in therapy, which represent a small proportion of the total recovered memory reports (Elliott, 1997; Wilsnack, Wonderlich, Kristjanson, Vogeltanz-Holm, & Wilsnack, 2002), were never corroborated in Geraerts small sample (n = 16).

Longitudinal studies also support the TM. Mechanic, Resick, and Griffin's (1998) study of amnesia postrape found that 37% of assaulted women reported some degree of amnesia at the 2-week point. At the 3-month marker, this number had dropped to 16%.

& Schacter, 1997), rather than immediately following the presentation of trauma-related words in the directed forgetting paradigm.

**Interidentity amnesia studies.** Interidentity amnesia in DID is a separate issue from that of dissociative amnesia in general. Authors from both TM and FM positions, including several of the authors of this review, have contributed to the general finding that implicit memories often cross dissociative identity barriers.

Interidentity amnesia has been studied as a paradigm for memory in DID since the late 19th and early 20th centuries (Prince & Peterson, 1908; see Dorahy, 2001). With renewed interest in multiple personality disorder and DID, this phenomenon has been examined to attempt to understand the nature of memory and amnesia in DID, often with contradictory findings (Eich, Macaulay, Loewenstein, & Dihle, 1997). In a series of studies designed to overcome these contradictions, Huntjens and others (Huntjens, 2003; Huntjens, Peters, Woertman, Van der Hart, & Postma, 2007) compared DID patients reporting mutually amnesic identities with simulator and normal controls. Studies included tests of neutral episodic information, perceptual and conceptual priming, procedural memory, transfer of trauma-related words, and stimulus valence as shown by affective priming. These researchers reported no objective evidence of interidentity amnesia in any of these studies. Huntjens (2003) concluded that dissociative amnesia in DID may have more to do with subjective appraisal and “metamemory” than actual lack of accessibility of memory between alternate identities.

Despite the amount of effort put into these studies, they have limitations. First, the notion of relatively stable, fixed “two-way” amnesic identities is based in the classical notion of DID as a small set of relatively unchanging, structured “personalities” with separate memory subsystems. This review is not the place to detail the TM-based view of the phenomenology of DID. Suffice it to say that the TM views DID as a posttraumatic developmental disorder with a relatively dynamic self-state system derived from a variety of developing intrapsychic, interpersonal, and psychosocial needs over time, and a phenomenology usually based in, overlap, interference, intrusion, and shifting (not simply switching) among personality states (Dell, 2006; Putnam, 1997). Further, this phenomenological model contrasts with the classical notion of well-defined identities with characteristics that can be reliably reproduced across clinical interviews and research trials (Dell, 2006; Putnam, 1997; Putnam, Zahn, & Post, 1990). Proponents of the TM—and, for that matter, proponents of the FM—do not take at face value DID identities’ prevalent beliefs that they actually are “real people” with varying demographic and psychological characteristics, including differing ages, genders, etc. Nor would proponents of either model take at face value other common beliefs that alternate identities are animals, mythical beings, internalized “outside” people, demons, or omniscient beings. Therefore, it is unclear why claims of two-way amnesia between identities should also be accepted at face value preferentially by either set of model theorists.

Thus, Forrest (1999, 2001), in a study of explicit memory in identities claiming coconsciousness, or shared memory, found evidence of interidentity amnesia, compared with normal and simulating controls, despite the identities’ beliefs in their coconsciousness. In additional support of the notion that alternate identities may not accurately assess their own subjective psychological characteristics, Loewenstein, Hamilton, Alagna, ReintalRe.5(LoleVruratsubjep)-247.5(yy)-408.8B5(Lol0.88897.6(clrest9.8(1997).)ders.)-3e



Table 5  
Review of Psychobiological Studies of Dissociation

Study	Sample description	Measures and method	Measures of dissociation and diagnosis	Results of interest in brief
Becker-Blease et al., 2004	75 unrelated adoptive sibling pairs; 218 MZ twins, and 173 DZ twins	Heritability and genetics Twin study of $r^2$ of trait dissociation; parent and teacher rated trait dissociation	6 trait dissociative items from CBCL	$h^2$ of dissociation = .60, $c^2$ = .00
Jang et al., 1998	General population sample of 177 MZ twins and 152 DZ twins	Twin study of $r^2$ of trait dissociation	DES-T, DES	$r^2$ of DES-T and DES = .48 and .55, respectively; $c^2$ = .00 for both DES-T and DES
Lochner et al., 2007	83 OCD participants	Genetic study of 5-HTTLPR, childhood trauma history and trait dissociation	DES-T, DES, CTQ	Childhood trauma and 5-HTT genotype predicted 22% of the variance in DES-T scores. Moderate correlations between CTQ and DES-T scores with SS genotype; association nonsignificant with LL genotype.
Pleper et al., 2011	184 twin pairs	Twin study of $r^2$ of trait dissociation, 5-HTTLPR, trauma history and trait dissociation	DES-T, DES	$r^2$ of DES-T and DES = .43 and .44, respectively; $c^2$ = .00 for both DES-T and DES. Participants with the SS genotype who also had high depressive symptoms and trauma had highest DES-T scores.
Savitz et al., 2008	178 individuals from 35 families bipolar proband and one additional first-degree relative with bipolar disorder	Study of genes related to COMT, polymorphism, trauma and trait dissociation	DES	DES scores predicted by the interaction of COMT genotype with childhood trauma; DES scores highest in individuals with the Val/Val genotype with childhood trauma
Tellegen et al., 1988	217 MZ and 114 DZ adult twins reared together and 44 MZ and 27 DZ adult twins reared apart	Twin study of $r^2$ of trait absorption	MPQ	$r^2$ of absorption = .50, $c^2$ = .03
Waller & Ross, 1997	280 MZ and 148 DZ twins	Twin study of $r^2$ of trait dissociation	DES-T	$r^2$ of DES-T scores = .00; $c^2$ = .45
Bonanno et al., 2003	103 women, 48 with documented CSA	HR while participants spoke of the "most distressing event" was contrasted with baseline HR	Psychophysiology (trait dissociation) ADES-T	ADES-T scores correlated negatively with increases in HR ( $r$ = -.24) and facial expressions ( $r$ = -.21) during discussion of distressing events (relative to baseline). DES correlated with SCRs to the video (.34); fantasy proneness showed null effects (.18, ns).
Giesbrecht et al., 2007	62 undergraduates	Viewed a provocative video while HR was measured	SOBS	Within trauma groups, higher DES (-.24) and DSS ( $r$ = -.20) related with lower HRV, whereas PDEQ was not correlated with either.
Hauschildt et al., 2011	26 trauma exposed with PTSD, 26 trauma exposed without PTSD, 18 nontrauma controls	HRV recorded during videos of varying emotional valence	DES, DSS, PDEQ	Lower HR was associated with higher derealization ( $r$ = -.29) and higher identity alteration ( $r$ = -.33).
Koopman et al., 2004	41 delinquent adolescents	Randomly assigned to either talk about their most stressful life experience or talk freely while HR was measured	ABD-D	
Sierra et al., 2002	15 DD patients, 15 HC, 11 anxiety controls			

Table 5 (continued)

Study	Sample description	Measures and method	Measures of dissociation and diagnosis	Results of interest in brief
Neuroendocrinology				
State dissociation Morgan et al., 2001	44 healthy male soldiers	NE, EPI, NPY, and plasma/salivary cortisol assessed before, during, and after exposure to physical and mental stress	CADSS	Increased CADSS associated with decreased cortisol during stress ( $r = -.49$ ) and increased cortisol 24 hr subsequently ( $r = -.46$ )
Trait dissociation Koopman et al., 2003	49 women with PTSD related to CSA	Five salivary cortisol samples collected at the beginning, end, and 1, 24, and 48 hr after an interview about stress and childhood trauma	SASRQ	High dissociators had elevated salivary cortisol 24 hr after the interview.
Schechter et al., 2004	41 mothers of young children with PTSD from interpersonal trauma	Maternal perception of children and behavior assessed during separation-reunion sequences. Salivary cortisol collected from mothers before and 30 min after reunions	HADS-I	Baseline salivary cortisol negatively correlated with severity of trait dissociation ( $r = -.31$ ), but cortisol reactivity to separation-reunion was nonsignificant correlated with severity of dissociation ( $r = .15, ns$ )
Simeon et al., 2007	46 DD without PTSD, 35 PTSD, 58 HC	24-hr urine and serial blood samples collected before and after DST and TSST	DES	DD had higher basal cortisol in urine (but not plasma) compared with HC. DD group had greater resistance to and faster escape from DST. No differences in cortisol reactivity. DES correlated negatively with peak cortisol reactivity to the TSST ( $r = -.43$ ).
Simeon et al., 2008	21 high exposure and 10 nontrauma HC without major exposure to the World Trade Center attack	24-hr urine cortisol after DST. During TSST, plasma cortisol changes, HR, and BP assessed during rest and at peak response	DES	DES negatively correlated with plasma cortisol levels at 08.00 h post-DST ( $r = -.56$ ), but not with baseline urinary cortisol ( $r = -.29, ns$ ), DST suppression ( $r = .12$ ), or cortisol reactivity to the TSST ( $r = -.18$ ). DES negatively correlated with resting systolic BP ( $r = -.54$ ) and peak HR during the TSST ( $r = -.48$ ) but unrelated to other BP and HR measures during rest and TSST.
Neuroimaging				
Structural imaging Irlle et al., 2009	10 PTSD with either DA or DID, 25 HC	MRI of total brain volume, bilateral amygdala, and bilateral hippocampus	SCID-D diagnosed DID and DA	Volumes of left 31% and right 29% amygdala and left 17% and right 11.0% hippocampal volumes were reduced when compared with HCs, but correlated with PTSD symptom severity rather than DA/DID symptoms.
Vermetten et al., 2006	15 DID, 23 HC	MRI determined hippocampal and amygdala volume	SCID-D diagnosed DID	Hippocampal volume 19% less in DID but confounded by age differences. Amygdala volume 32% less in DID, but only the effect of right amygdala volume still significant after covarying age.
Weniger et al., 2008	13 DID or DA, 25 HC, 10 PTSD	MRI scan of amygdala and hippocampal size	SCID-D diagnosed DID and DA	Neither amygdala nor hippocampal volumes differed between the DID/DA group and HC.
Functional imaging Brand et al., 2009	14 DA, 19 HC	PET scan acquired during eyes-closed resting state	DA diagnosis made according to DSM-IV	DA participants showed less metabolism in the right inferolateral PFC.
Eizinga et al., 2007	16 DID or DDNOS, 16 HC	fMRI scanning during verbal test of working memory	DES	Working memory nonsignificant. DID/DDNOS greater response as a function of increasing task difficulty, relative to HC, within left anterior PFC, DLPFC, and parietal lobe (BA 40).

(table continues)

Table 5 (continued)

Study	Sample description	Measures and method	Measures of dissociation and diagnosis	Results of interest in brief
Felmingham et al., 2008	12 PTSD displaying dissociative reactions, 11 PTSD who did not dissociate			

Table 5 (continued)

Study	Sample description	Measures and method	Measures of dissociation and diagnosis	Results of interest in brief



trauma exposure and response to both idiographic and standardized stimuli (e.g., McTeague et al., 2010).

A growing number of studies have examined cortisol response as a measure of stress reactivity and functioning of the hypothalamic–pituitary–adrenal axis in individuals as a function of dissociative symptoms. These studies support the basic principle of the TM that traumatic stress plays a causal role in dissociative symptoms. However, most studies to date have examined peritraumatic dissociation only (e.g., Ladwig et al., 2002; Neylan et al., 2005; Nixon et al., 2005). Higher (Simeon et al., 2007), lower (Schechter et al., 2004), and null effects have been observed for basal cortisol in comparisons of individuals high versus low in dissociative symptoms. Cortisol reactivity to psychological stressors was decreased in response to combat training as a function of state dissociation (Morgan et al., 2001), but Simeon et al. (2007, 2008) did not find decreased (or increased) cortisol reactivity in response to the Trier Social Stress Test. Finally, Koopman et al. (2003) observed increased salivary cortisol in individuals reporting greater trait dissociative symptoms (see Table 5). Phillips et al. (2001) observed less difference in only 1 day (but not immediately or 2 days) after being interviewed about traumatic life events. Discrepant findings across studies may suggest that patterns of arousal differentiating high and low dissociators within PTSD groups may change over time. In addition, future studies of cortisol reactivity to psychological stressors as a function of trait dissociation should examine the extent that individuals experience state dissociation in response to the stressor.

The significance of documented psychophysiological and neuroendocrine correlates of self-reported dissociative symptoms can be interpreted from either the FM or the TM perspective. FM theorists can maintain that objective psychophysiological responses to stimuli reminiscent of trauma may represent solely the individuals' belief that they have experienced trauma, a belief that may be unfounded in reality. For example, McNally et al. (2004) found that heart rate, SCR, and left frontal electromyography was also observed in PTSD patients reporting dissociative symptoms increased more significantly in individuals believing themselves to be alien abductees than in comparison volunteers when exposed to auditory recounting of alien abduction experiences. However, McNally et al. did not distinguish between high and low dissociation groups in their analysis.

In comparison, TM theorists may note that behavioral and psychophysiological responses observed in reportedly traumatized dissociative subjects closely match those often observed in animals within the context of inescapable predatory threat, a behavioral pattern referred to in the animal literature as tonic immobility (Bracha & Maser, 2008; Bracha, Ralston, Matsukawa, Williams, & Bracha, 2004; Marx, Forsyth, Gallup, Fusch Lexington, 2008; Moskowitz, 2004). Within the state of tonic immobility, an animal takes upon itself an outwardly passive defensive response involving inhibition of movement, muscular rigidity or limpness, and evidently unfocused concentration (e.g., unfocused gaze, eye closture), a behavioral and psychophysiological state that has been associated with increased analgesia. These characteristics bear a resemblance to certain dissociative states as discussed above (Frewen & Lanius, 2006; Nijenhuis, Vanderlinden, & Spinhoven, 1998). Tonic immobility to date has been examined primarily in its relevance to trauma and PTSD as opposed to dissociative symptoms specifically, although researchers have discussed its particular relevance to dissociative symptoms in PTSD (Bovin, Jager-Hyman, Gold, Marx, & Sloan, 2008; Fiszman et al., 2008; Heidt, Marx, & Forsyth, 2005; Humphreys, Sauder, Martin, & Marx, 2010; Rocha-Rego et al., 2009). Further-

more, psychometrically measured tonic immobility correlates with dissociative symptoms (Abrams, Carleton, Taylor, & Asmundson, 2009). In short, the animal literature on tonic immobility affords a translational model informing the psychophysiological study of dissociative symptoms. These studies support the basic principle of the TM that traumatic stress plays a causal role in dissociative symptoms. Higher (Simeon et al., 2007), lower (Schechter et al., 2004), and null effects have been observed for basal cortisol in comparisons of individuals high versus low in dissociative symptoms. Cortisol reactivity to psychological stressors was decreased in response to combat training as a function of state dissociation (Morgan et al., 2001), but Simeon et al. (2007, 2008) did not find decreased (or increased) cortisol reactivity in response to the Trier Social Stress Test. Finally, Koopman et al. (2003) observed increased salivary cortisol in individuals reporting greater trait dissociative symptoms (see Table 5). Phillips et al. (2001) observed less difference in only 1 day (but not immediately or 2 days) after being interviewed about traumatic life events. Discrepant findings across studies may suggest that patterns of arousal differentiating high and low dissociators within PTSD groups may change over time. In addition, future studies of cortisol reactivity to psychological stressors as a function of trait dissociation should examine the extent that individuals experience state dissociation in response to the stressor.

Neuroimaging studies have examined emotional processing in subjects with DPD, and trauma memory and/or pain processing in individuals with PTSD or borderline personality disorder (BPD) with prominent dissociative symptoms, with a common finding being either increased or decreased response in medial prefrontal cortex and limbic regions accompanying dissociative symptoms (see Table 5). Phillips et al. (2001) observed less difference in emotional processing regions of the brain, most notably the insula, and a greater frontal response, in people with DPD when viewing disvalenced photographs. Among PTSD patients, individuals exhibiting state depersonalization in response to trauma reminders also showed an increased response within midline anterior regions including the dorsal anterior cingulate cortex and the medial prefrontal cortex (Hopper, Frewen, Sack, Lanius, & Van der Kolk, 2007; Lanius et al., 2010, 2005, 2002). In comparison, null effects were observed in 10 participants with DPD, although during a subsequent recognition test for emotional words, healthy controls activated the medial prefrontal cortex more so than individuals with DPD (Medford et al., 2006). Less response within medial prefrontal cortex was also observed in PTSD patients reporting dissociative symptoms in response to threatening facial expressions (Felmingham et al., 2008). Increased midcingulate and insula response in patients with BPD and comorbid PTSD was observed in conjunction with reduced pain sensitivity during script-induced dissociative states (Ludäscher et al., 2010). Thus, functional neuroimaging studies increasingly implicate the anterior cingulate and limbic basis for positive symptoms of dissociative disorders and dissociative symptomatology, most notably those of depersonalization and analgesia. Recently, neuroimaging studies have also sought to investigate the basis of negative symptoms of dissociation, including dissociative amnesia and interictal amnesia. Findings in 14 individuals with dissociative amnesia and comorbid PTSD showed decreased metabolism within the right inferolateral prefrontal cortex typically within the right hemisphere, in individuals whose amnesia was documented to have been provoked by traumatic and/or stressful events (review by Staniloiu & Markowitsch, 2010; see also Staniloiu, Markowitsch, & Brand, 2010). Vermetten, Schmahl, Lindner, Loewenstein, and Bremner (2006) observed reduced volume of the hippocampus and amygdala in individuals with DID. This result was not replicated in a subsequent study, where brain morphological changes were reported to be associated with a PTSD diagnosis, not with a dissociative disorder diagnosis without PTSD (Irlle, Lange, Sachsse, & Weniger, 2009; Weniger, Lange, Sachsse, & Irlle, 2008). However, in

these latter studies, only four of 13 trauma-exposed individuals met SCID-D diagnostic criteria for DID. Most met diagnostic criteria for dissociative amnesia, and no data are reported on which dissociative patients met diagnostic criteria for PTSD. Accordingly, further studies will be needed to more completely elucidate whether the Vermetten et al. findings can be better explained by comorbid PTSD, by DID, or by both disorders.

The reviewed neuroimaging studies were not designed to address the present question regarding the degree to

that each of these additions to the TM are proposed as moderators of the dissociation–trauma relationship, not as mediators of the relationship. Future researchers would be better served by designs that include relevant variables as independent grouping factors (e.g., intrafamilial vs. extrafamilial abuse, low vs. high family pathology) so that simple effects and interactions can be examined.

Recent studies have begun to answer this question. Data from the National Comorbidity Study–Replication report that multiple forms of childhood adversity, including childhood maltreatment and family dysfunction, covary strongly together, such that it may not be possible to separate the effects of maltreatment from a pathogenic family environment in which multiple forms of neglect and abuse occur (Green et al., 2010; McLaughlin et al. 2010; Scott, Varghese, & McGrath, 2010). Trickett et al. (2011) came to a similar conclusion in their review of the many pathological outcomes of childhood sexual abuse, including increased dissociation. These adverse outcomes are difficult to completely parcel out from the manifold harms caused by the pathogenic family environment in which childhood sexual abuse, physical abuse, emotional abuse, and neglect occur.

### **Should We Discount the Nonobjective Trauma Studies?**

Regarding the issue of objective and subjective measures of trauma in general, it is certainly true that much research on trauma is conducted with participants whose traumatic background has not been independently verified. This, however, is the norm rather than the exception in most areas of psychology. In comparing nonsmokers with light and heavy smokers on rates of varying diseases, seldom are there external documents verifying the number of cigarettes per week actually consumed. Salivary cotinine levels have been used to document abstinence after intervention, but are used less now because of the high correspondence between these levels and self-report (Yeager & Krosnick, 2010). The number of bingeing or purging episodes for the bulimic are virtually never verified, nor is there an objective verification that the fantasy-prone individual actually spends more time fantasizing. Thus, in a wide range of fields, it is understood that self-report contains measurement error, and independent studies are conducted to show that the criterion-positive group (e.g., alcoholic, sexually abused, bulimic) is reliably more likely to contain criterion-positive individuals than the self-reported criterion-negative group.

Unfortunately, longitudinal studies cannot provide a full answer to the question here, since the individual who first reports sexual abuse as an adult cannot dependably be labeled as a false report (even if the same individual denied it as a child), because alternative hypotheses of shame or fear serving to silence the child from disclosing abuse are viable possibilities. Twenty-year follow-ups of a large sample of abused children and matched controls revealed large omission rates for those asked if they had experienced prior physical abuse (38%–40% in Widom & Shepard, 1996) and prior

time should be slow or nonexistent and unrelated to trauma or trauma treatment. Instead, as the TM suggests, dissociation drops over the course of the 1st year after trauma for most individuals (e.g., Feeny, Zoellner, Fitzgibbons, & Foa, 2000; Feeny, Zoellner,

## **Psychobiology of Dissociation as a Regulatory Response to Trauma**

Extant research supports the TM of dissociation as a regulatory response to fear or other extreme emotion with measurable biological correlates. The strong caveat here is that, to our knowledge, most research has not been done with FM and TM theories in mind, and thus has not included measures of fantasy proneness or suggestibility. Nonetheless, biological researchers have found trauma-related theories (e.g., tonic immobility) to be useful in synthesizing findings from animal and human samples. Compelling alternative heuristics that are not trauma related have yet to appear.

### **Summary**

Finally, in future research, we recommend the careful analysis of varying alternative causal models; attempts to differentiate mediators, moderators, and risk factors; the avoidance of use of outlier studies to make theoretical arguments; and attention to measurement issues in all conceptual areas (dissociation, fantasy proneness and false memory) to further this complicated and fascinating dialogue. Our review of current research suggests that trauma and dissociation are connected for psychological and neurobiological reasons, and fantasy proneness is not the explanation.

### **References**

- Abramowitz, E. G., & Lichtenberg, P. (2010). A new hypnotic technique for treating combat-related posttraumatic stress disorder: A prospective open study. *International Journal of Clinical and Experimental Hypnosis*, 58, 316–328. doi:10.1080/00207141003760926
- Abrams, M. P., Carleton, R. N., Taylor, S., & Asmundson, G. J. G. (2009).



- disorders. *Psychotherapy and Psychosomatics*, 76, 400–401. doi: 10.1159/000107570
- Dalenberg, C. J., & Paulson, K. (2009). The case for the study of “normal” dissociation processes. In P. F. Dell & J. A. O’Neil (Eds.), *Dissociation and the dissociative disorder DSM–V and beyond* (pp. 145–154). New York, NY: Routledge.
- Deese, J. (1959). On the prediction of occurrence of particular verbal intrusions in immediate recall. *Journal of Experimental Psychology*, 58, 17–22. doi:10.1037/h0046671
- Dell, P. F. (2006). The Multidimensional Inventory of Dissociation (MID): A comprehensive measure of pathological dissociation. *Journal of Trauma & Dissociation*, 7, 77–106. doi:10.1300/J229v07n02\_06
- DePrince, A. P., Allard, C. B., Oh, H., & Freyd, J. J. (2004). What’s in a name for memory errors? Implications for ethical issues arising from the use of the term “false memory” for errors in memory for details. *Law, Ethics & Behavior*, 14, 201–233. doi:10.1207/s15327019eb1403\_1
- DePrince, A. P., & Freyd, J. J. (2001). Memory and dissociative tendencies: The roles of attentional context and word meaning in a directed forgetting task. *Journal of Trauma & Dissociation*, 2, 67–82. doi: 10.1300/J229v02n02\_06
- DePrince, A. P., & Freyd, J. J. (2004). Forgetting trauma stimuli. *Psychological Science*, 15, 488–492. doi:10.1111/j.0956-7976.2004.00706.x
- DePrince, A. P., Weinzierl, K. M., & Combs, M. D. (2008). Stroop performance, dissociation, and trauma exposure in a community sample of children. *Journal of Trauma & Dissociation*, 9, 209–223. doi: 10.1080/15299730802048603
- de Ruiter, M. B., Elzinga, B. M., & Phaf, R. H. (2006). Dissociation: Cognitive capacity or dysfunction? *Journal of Trauma & Dissociation*, 7, 115–134. doi:10.1300/J229v07n04\_07
- de Ruiter, M. B., Phaf, R. H., Elzinga, B. M., & Van Dyck, R. (2004). Dissociative style and individual differences in verbal working memory span. *Consciousness and Cognition*, 13, 821–828. doi:10.1016/j.concog.2004.08.002
- Devilley, G. J., Ciociari, J., Piesse, A., Sherwell, S., Zammit, S., Cook, F., & Turton, C. (2007). Dissociative tendencies and memory performance on directed forgetting task. *Psychological Science*, 18, 212–217. doi: 10.1111/j.1467-9280.2007.01875.x
- De Wachter, D., Lange, A., Vanderlinden, J., Pouw, J., & Strubbe, E. (2006). The influence of current stress on dissociative experiences: An exploratory study in a non-clinical population. *Journal of Trauma & Dissociation*, 7, 87–96. doi:10.1300/J229v07n01\_07
- Diseth, T. H. (2006). Dissociation following traumatic medical treatment

Feeny, N. C., Zoellner, L. A., Fitzgibbons, L. A., & Foa, E. B. (2000).  
Exploring the roles of emotional numbing, depression, and dissociation  
in PTSD. *Journal of Traumatic Stress*, 13, 489–498. doi:10.1023/A:  
1007789409330



variability in response to affective scenes in posttraumatic stress disorder. *Biological Psychology*, 88, 215–222. doi:10.1016/j.biopsycho.2011.08.004

Hébert, M., & Tourigny, M. (2010). Effects of a psychoeducational group

- disorders? An experimental investigation. *Behavioral and Cognitive Psychotherapy*, 36, 221–234. doi:10.1017/S1352465807004080
- Klinger, E., Henning, V. R., & Janssen, J. M. (2009). Fantasy-proneness dimensionalized: Dissociative component is related to psychopathology, daydreaming as such is not. *Journal of Research in Personality*, 43, 506–510. doi:10.1016/j.jrp.2008.12.017
- Koopman, C., Carrion, V., Butler, L. D., Sudhakar, S., Palmer, L., & Steiner, H. (2004). Relationships of dissociation and childhood abuse and neglect with heart rate in delinquent adolescents. *Journal of Traumatic Stress*, 17, 47–54. doi:10.1023/B:JOTS.0000014676.83722.35
- Koopman, C., Sephton, S., Abercrombie, H. C., Classen, C., Butler, L. D., Gore-Felton, C., . . . Spiegel, D. (2003). Dissociative symptoms and cortisol responses to recounting traumatic experiences among childhood sexual abuse survivors with PTSD. *Journal of Trauma & Dissociation*, 4, 29–46. doi:10.1300/J229v04n04\_03
- Koutstaal, W., & Schacter, D. L. (1997). Intentional forgetting and voluntary thought suppression: Two potential methods for coping with childhood trauma. In L. J. Dickstein, M. B. Riba, & J. M. Oldham (Eds.), *Review of psychiatry* (Vol. 16, pp. 117–121). Washington, DC: American Psychiatric Press.
- Kraemer, H. C., Stice, E., Kazdin, A., Offord, D., & Kupfer, D. (2001). How do risk factors work together? Mediators, moderators, and independent, overlapping, and proxy risk factors. *American Journal of Psychiatry*, 158, 848–856. doi:10.1176/appi.ajp.158.6.848
- Ladwig, K.-H., Marten-Mittag, B., Deisenhofer, I., Hofmann, B., Schaperer, J., Weyerbrock, S., . . . Schmitt, C. (2002). Psychophysiological correlates of peritraumatic dissociative responses in survivors of life-threatening cardiac events. *Psychopathology*, 35, 241–248. doi:10.1159/000063825
- Lanius, R. A., Vermetten, E., Loewenstein, R. J., Brand, B., Schmahl, C., Bremner, J. D., & Spiegel, D. (2010). Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *American Journal of Psychiatry*, 167, 640–647. doi:10.1176/appi.ajp.2009.09081168
- Lanius, R. A., Williamson, P. C., Bluhm, R. L., Densmore, M., Boksman, K., Neufeld, R. W. J., . . . Menon, R. S. (2005). Functional connectivity of dissociative response in posttraumatic stress disorder: A functional magnetic resonance imaging investigation. *Biological Psychiatry*, 57, 873–884. doi:10.1016/j.biopsych.2005.01.011
- Lanius, R. A., Williamson, P. C., Boksman, K., Densmore, M., Gupta, M., Neufeld, R. W. J., . . . Menon, R. S. (2002). Brain activation during script-driven imagery induced dissociative responses in PTSD: A functional magnetic resonance imaging investigation. *Biological Psychiatry*, 52, 305–311. doi:10.1016/S0006-3223(02)01367-7
- Leavitt, F. (1997). False attribution of suggestibility to explain recovered memory of childhood sexual abuse following extended amnesia. *Child Abuse & Neglect*, 21, 265–272. doi:10.1016/S0145-2134(96)00171-8
- Leavitt, F. (1999). Dissociative experiences scale taxon and measurement of dissociative pathology: Does the taxon add to an understanding of dissociation and its associated pathologies? *Journal of Clinical Psychology in Medical Settings*, 6, 427–440. doi:10.1023/A:1026275916184
- Leavitt, F. (2001). MMPI profile characteristics of women with varying levels of normal dissociation. *Journal of Clinical Psychology*, 57, 1469–1477. doi:10.1002/jclp.1110
- Leonard, K. N., Telch, M. J., & Harrington, P. J. (1999). Dissociation in the laboratory: A comparison of strategies. *Behaviour Research and Therapy*, 37, 49–61. doi:10.1016/S0005-7967(98)00072-2
- Levin, R., Sirof, B., Simeon, D., & Guralnick, O. (2004). Role of fantasy proneness, imaginative involvement, and psychological absorption in depersonalization disorder. *Journal of Nervous and Mental Disease*, 192, 69–71. doi:10.1097/01.nmd.0000106003.46153.54
- Levin, R., & Spei, E. (2004). Relationship of purported measures of pathological and nonpathological dissociation to self-reported psychological distress and fantasy immersion. *Assessment*, 11, 160–168. doi:10.1177/1073191103256377
- Lev-Wiesel, R., Daphna-Tekoah, S., & Hallak, M. (2009). Childhood sexual abuse as a predictor of birth-related posttraumatic stress and postpartum posttraumatic stress. *Child Abuse & Neglect*, 33, 877–887. doi:10.1016/j.chiabu.2009.05.004
- Lewis, D. O., Yeager, C. A., Swica, Y., Pincus, J. H., & Lewis, M. (1997). Objective documentation of child abuse and dissociation in 12 murderers with dissociative identity disorder. *American Journal of Psychiatry*, 154, 1703–1710.
- Lilienfeld, S. O., Kirsch, I., Sarbin, T. R., Lynn, S. J., Chaves, J. F., Ganaway, G. K., & Powell, R. A. (1999). Dissociative identity disorder and the sociocognitive model: Recalling the lessons of the *Psychological Bulletin*, 125, 507–523. doi:10.1037/0033-2909.125.5.507
- Lindsay, S., & Briere, J. (1997). The controversy regarding recovered memories of childhood sexual abuse: Pitfalls, bridges, and future directions. *Journal of Interpersonal Violence*, 12, 631–647. doi:10.1177/088626097012005002
- Little, L. (1996). Source monitoring and dissociative experiences in DID and non-DID inpatients: A multicenter study. (Doctoral dissertation). Available from ProQuest Dissertations and Theses database. (UMI No. 9805139)
- Lochner, C., Seedat, S., Hemmings, S. M., Kinnear, C. J., Corfield, V. A., Niehaus, D. J. H., . . . Stein, D. J. (2004). Dissociative experiences in obsessive-compulsive disorder and trichotillomania: Clinical and genetic findings. *Comprehensive Psychiatry*, 43, 384–391. doi:10.1016/j.comppsy.2004.03.010
- Lochner, C., Seedat, S., Hemmings, S. M. J., Moolman-Smook, J. C., Kidd, M., & Stein, D. J. (2007). Investigating the possible effects of trauma experiences and HTT on the dissociative experiences of patients with OCD using path analysis and multiple regression. *Neuropsychobiology*, 56, 6–13. doi:10.1159/000109971
- Loewenstein, R. J., Hamilton, J., Alagna, S., Reid, N., & deVries, M. (1987). Experiential sampling in the study of multiple personality disorder. *American Journal of Psychiatry*, 144, 9–24.
- Loftus, E., & Ketcham, K. (1994). *The myth of repressed memory: False memories and allegations of sexual abuse*. New York, NY: St. Martin's Press.
- Ludäscher, P., Bolus, M., Lieb, K., Phillipsen, A., Jochims, A., & Schmahl, C. (2007). Elevated pain thresholds correlate with dissociation and aversive arousal in patients with borderline personality disorder. *Psychiatry Research*, 149, 291–296. doi:10.1016/j.psychres.2005.04.009
- Ludäscher, P., Valerius, G., Stigmayer, C., Mauchnik, J., Lanius, R. A., Bohus, M., & Schmahl, C. (2010). Pain sensitivity and neural processing during dissociative stress in patients with borderline personality disorder with and without comorbid posttraumatic stress disorder: A pilot study. *Journal of Psychiatry and Neuroscience*, 35, 177–184. doi:10.1503/jpn.090022
- Lynn, S. J., & Rhue, J. W. (1986). The fantasy-prone person: Hypnosis, imagination, and creativity. *Journal of Personality and Social Psychology*, 51, 404–408. doi:10.1037/0022-3514.51.2.404
- Lyons-Ruth, K., Dutra, L., Schuder, M. R., & Bianchi, I. (2006). From infant attachment disorganization to adult dissociation: Relational adaptations or traumatic experiences? *Psychiatric Clinics of North America*, 29, 63–86. doi:10.1016/j.psc.2005.10.011
- Lyttle, N., Dorahy, M. J., Hanna, D., & Huntjens, R. J. C. (2010). Conceptual and perceptual priming and dissociation in chronic posttraumatic stress disorder. *Journal of Abnormal Psychology*, 119, 777–790. doi:10.1037/a0020894
- Macfie, J., Cicchetti, D., & Toth, S. L. (2001a). The development of dissociation in maltreated preschool-aged children. *Development and Psychopathology*, 13, 233–254. doi:10.1017/S0954579401002036
- Macfie, J., Cicchetti, D., & Toth, S. L. (2001b). Dissociation in maltreated

versus nonmaltreated preschool-aged children *Child Abuse & Neglect*,  
25, 1253–1267. doi:10.1016/S0145-2134(01)00266-6



childhood: True and false reports. *Journal of Experimental Psychology: Applied*, 14, 373–391. doi:10.1037/a0014309

Rauschenberger, S. L., & Lynn, S. J. (1995). Fantasy proneness, DSM-III-R Axis I psychopathology, and dissociation. *Journal of Abnormal Psychology*, 104, 373–380. doi:10.1037/0021-843X.104.2.373

Rauschenberger, S. L., & Lynn, S. J. (2002–2003). Fantasy-proneness, negative affect, and psychopathology. *Imagination, Cognition and Personality*, 22, 239–255. doi:10.2190/AEXB-Y3JR-Y7T2-s0A7R

Reinders, A. A. T. S., Nijenhuis, E. R. R. Reinders, A. M. (2006). Fantasy-proneness, dissociation, and psychopathology. *Journal of Experimental Psychology: Applied*, 12, 1–12. doi:10.1037/1076-898X.12.1.1

(Eds.), *Dissociation and the dissociative disorders: DSM-V and beyond* (pp. 435–444). New York, NY: Routledge.

Simeon, D., Guralnik, O., Hazlett, E. A., Spiegel-Cohen, J., Hollander, E., & Buchsbaum, M. S. (2000). Feeling unreal: A PET study of depersonalization disorder. *American Journal of Psychiatry*, *157*, 1782–1788. doi:10.1176/appi.ajp.157.11.1782

Simeon, D., Knutelska, M., Yehuda, R., Putnam, F., Schmeidler, J., & Smith, L. M. (2007). Hypothalamic–pituitary–adrenal axis function in dissociative disorders, PTSD, and healthy volunteers. *Biological Psychiatry*, *61*,

- PTSD patients' narratives during prolonged exposure therapy: A replication and extension *Journal of Traumatic Stress*, 12, 255–258. doi: 10.1023/A:1015263513654
- Vasterling, J. J., & Brewin, C. R. (Eds.). (2005). *Neurobiology of PTSD: Biological, cognitive, and clinical perspectives*. New York, NY: Guilford Press.
- Vasterling, J. J., Duke, L. M., Brailey, K., Constans, J. I., Allain, A. N., & Sutker, P. B. (2002). Attention, learning, and memory performances and intellectual resources in Vietnam veterans: PTSD and no disorder comparisons. *Neuropsychology*, 16, 1–14. doi:10.1037/0894-4105.16.1.5
- Veltman, D. J., de Ruiter, M. B., Rombouts, S. A. R. B., Lazeron, R. H. C., Barkhof, F., Van Dyck, R., . . . Phaf, R. H. (2005). Neurophysiological correlates of increased verbal working memory in high-dissociative participants: A functional MRI study. *Psychological Medicine*, 35, 175–185. doi:10.1017/S0033291704002971
- Vermetten, E., Schmahl, C., Lindner, S., Loewenstein, R. J., & Bremner, J. D. (2006). Hippocampal and amygdalar volumes in dissociative identity disorder. *American Journal of Psychiatry*, 163, 630–636. doi: 10.1176/appi.ajp.163.4.630
- Waldo, T. G., & Merritt, R. D. (2000). Fantasy proneness, dissociation, and DSM-IV Axis II symptomatology. *Journal of Abnormal Psychology*, 109, 555–558. doi:10.1037/0021-843X.109.3.555
- Waller, N. G., Putnam, F. W., & Carlson, E. B. (1996). Types of dissociation and dissociative types: A taxometric analysis of dissociative experiences. *Psychological Methods*, 1, 300–321. doi:10.1037/1082-989X.1.3.300
- Waller, N. G., & Ross, C. A. (1997). The prevalence and biometric structure of pathological dissociation in the general population: Taxometric and behavior genetic findings. *Journal of Abnormal Psychology*, 106, 499–510. doi:10.1037/0021-843X.106.4.499
- Watson, D. (2003). Investigating the construct validity of the dissociative taxon: Stability analyses of normal and pathological dissociation. *Journal of Abnormal Psychology*, 112, 298–305. doi:10.1037/0021-843X.112.2.298
- Watson, D., Wu, K. D., & Cutshall, C. (2004). Symptom subtypes of obsessive-compulsive disorder and their relation to dissociation. *Anxiety Disorders*, 18, 435–458. doi:10.1016/S0887-6185(03)00029-X
- Weniger, G., Lange, C., Sachsse, U., & Irle, E. (2008). Amygdala and hippocampal volumes and cognition in adult survivors of childhood abuse with dissociative disorder. *Acta Psychiatrica Scandinavica*, 118, 281–290. doi:10.1111/j.1600-0447.2008.01246.x
- Widom, C. S., & Morris, S. (1997). Accuracy of adult recollections of childhood victimization: Part 2. Childhood sexual abuse. *Psychological Assessment*, 9, 44–46. doi:10.1037/1040-3590.9.1.34
- Widom, C. S., & Shepard, R. L. (1996). Accuracy of adult recollections of childhood victimization: Part 1. Childhood physical abuse. *Psychological Assessment*, 8, 412–421. doi:10.1037/1040-3590.8.4.412